# DRUG WARNING

High dose MDMA (ecstasy) tablets recently found in NSW – also containing a synthetic cathinone

### **Know the risks**

- Synthetic cathinones are stimulants that may have similar effects to MDMA and other amphetamines.
   There is limited information on the specific effects of dipentylone.
- MDMA tablets are often poorly manufactured and the amount of MDMA and other substances in a tablet can vary a lot, even within the same batch.
- You are at increased risk of harm if you:
  - Take multiple MDMA (ecstasy) tablets or capsules
  - Take a higher-dose MDMA tablet or capsule
  - Take MDMA in a hot environment like a music festival or indoor dance party
- The time from taking the drug to feeling any effect can vary between people and tablets. If it's not working, don't re-dose.
- High-dose MDMA use has been linked to death and other serious harms.

#### Effects to look out for

- · Feeling really hot and sweaty
- Racing heart/pulse or light-headedness
- Rigid muscles (e.g. difficulty walking), uncontrolled repetitive movements, seizures
- Confusion or agitation, feeling aggressive, difficult to rouse or unconscious
- Vomiting

If you or others experience these symptoms after taking MDMA, get help immediately.

## **Getting help**

If you or your friends see the warning signs of overdose:

Seek help immediately from your nearest emergency department or call <u>Triple Zero (000)</u>. You won't get into trouble for seeking medical care.

Start CPR if someone is not responsive.

Skull shaped tablets with MYBRAND marking have been found containing high dose MDMA and a synthetic cathinone, dipentylone\*

The risk of harm is increased when MDMA is taken with other stimulants including synthetic cathinones, amphetamines or cocaine.

\* also known as N,N-dimethylpentylone



Pink-orange 182-185mg MDMA + dipentylone Blue 162mg MDMA + dipentylone Yellow 151mg MDMA

Other recently tested skull shaped tablets with MYBRAND marking have contained MDMA with ketamine or 2'-fluoro-2-oxo-PCE (a ketamine analogue, also known as 2F-NENDCK, or 2-fluoro-N-ethylnordeschloroketamine)

Tablet appearance is not a reliable indicator of drug contents

# Support and advice

For free and confidential advice:

- Call <u>Alcohol and Drug Information</u>
   <u>Service</u> (ADIS) on 1800 250 015 at any time 24/7.

  Start a <u>Web Chat</u> with an ADIS counsellor Mon-Fri, 8.30am-5pm.
- Call <u>NUAA</u> on 1800 644 413 (Mon-Fri 9am-5pm) to speak to a peer or visit <u>NUAA</u> for a range of resources.
- Call the <u>NSW Poisons Information Centre</u> on 13 11 26 for information on adverse effects from drugs.
- Visit <u>Your Room</u> for fact sheets and other resources.

For up to date alerts, visit <a href="https://www.health.nsw.gov.au/drugalerts">https://www.health.nsw.gov.au/drugalerts</a>

Issued 1 November 2024 © NSW Health.



